

## ILAE Academy and EpiCARE

### Bibliography to case study: Bibliography Marie 6 y/o (Sturge Weber)

Alkonyi B, Chugani HT, Juha C (2011) Transient focal cortical increase of interictal glucose metabolism in Sturge-Weber syndrome: Implications for epileptogenesis. *Epilepsia*. 52(7):1265–1272. doi: 10.1111/j.1528-1167.2011.03066.x

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3130832/pdf/nihms-276390.pdf>

Alkonyi B, Chugani HT, Karia S, Behen ME, Juhász C (2011) Clinical outcomes in bilateral Sturge-Weber syndrome. *Pediatr Neurol*. 2011 Jun;44(6):443-9. doi: 10.1016/j.pediatrneurol.2011.01.005.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3092166/pdf/nihms270217.pdf>

Arzimanoglou A, Andermann F, Aicardi J, Sainte-Rose C, Beaulieu MA, Villemure JG, Olivier A, Rasmussen T (2000) Sturge-Weber syndrome: indications and results of surgery in 20 patients. *Neurology*. Nov 28;55(10):1472-9. doi: 10.1212/wnl.55.10.1472.

Arzimanoglou A, Aicardi J (1992) The epilepsy of Sturge-Weber syndrome: clinical features and treatment in 23 patients. *Acta Neurol Scand. Suppl.* 140:18-22. doi: 10.1111/j.1600-0404.1992.tb04465.x.

Aupy J, Bonnet C, Arnould JS, Fernandez P, Marchal C, Zanotti-Fregonara P (2015) Focal inhibitory seizure with prolonged deficit in adult Sturge-Weber syndrome. *Epileptic Disord*. Sep;17(3):328-31.

doi: 10.1684/epd.2015.0755. [https://www.jle.com/download/epd-305194-29993-](https://www.jle.com/download/epd-305194-29993-focal_inhibitory_seizure_with_prolonged_deficit_in_adult_sturge_weber_syndrome-a.pdf)

[focal\\_inhibitory\\_seizure\\_with\\_prolonged\\_deficit\\_in\\_adult\\_sturge\\_weber\\_syndrome-a.pdf](https://www.jle.com/download/epd-305194-29993-focal_inhibitory_seizure_with_prolonged_deficit_in_adult_sturge_weber_syndrome-a.pdf)

Bar C, Kaminska A, Nabbout R (2018) Spikes might precede seizures and predict epilepsy in children with Sturge-Weber syndrome: A pilot study. *Epilepsy Res*. Jul;143:75-78. doi:

10.1016/j.eplesyres.2018.03.020.

Behen ME, Juhász C, Wolfe-Christensen C, Guy W, Halverson S, Rothermel R, Janisse J, Chugani JT (2011) Brain damage and IQ in unilateral Sturge-Weber syndrome: support for a "fresh start" hypothesis. *Epilepsy Behav*. Oct;22(2):352-7. doi: 10.1016/j.yebeh.2011.07.010.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3185171/pdf/nihms-314623.pdf>

Bernasconi A, Cendes F, Theodore WH, Gill RS, Koepp MJ, Hogan RE, Jackson GD, Federico P, Labate A, Vaudano AE, Blümcke I, Ryvlin P, Bernasconi N (2019) Recommendations for the use of structural magnetic resonance imaging in the care of patients with epilepsy: A consensus report from the International League Against Epilepsy Neuroimaging Task Force. *Epilepsia*. Jun;60(6):1054-1068. doi: 10.1111/epi.15612 <https://onlinelibrary.wiley.com/doi/epdf/10.1111/epi.15612>

Blumcke I, Spreafico R, Haaker G *et al* for the EEBB Consortium (2017) Histopathological Findings in Brain Tissue Obtained during Epilepsy Surgery. *N Engl J Med*. 377:1648-1656

doi : 10.1056/NEJMoa1703784.

<https://www.nejm.org/doi/pdf/10.1056/NEJMoa1703784?articleTools=true>

Bourgeois M, Crimmins DW, Santos de Oliveira R, Arzimanoglou A, Garnett M, Roujeau T, Di Rocco F, Sainte-Rose C (2007) Surgical treatment of epilepsy in Sturge-Weber syndrome in children. *J Neurosurg*. Jan;106(1 Suppl):20-8. doi: 10.3171/ped.2007.106.1.20.

Bye AM, Matheson JM, Mackenzie RA (1989) Epilepsy surgery in Sturge-Weber syndrome.

*Aust Paediatr J*. 1989 Apr;25(2):103-5. doi: 10.1111/j.1440-1754.1989.tb01428.x.

Comi A (2015) Current Therapeutic Options in Sturge-Weber Syndrome. *Semin Pediatr Neurol*.

Dec;22(4):295-301. doi: 10.1016/j.spen.2015.10.005.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4943027/pdf/nihms789047.pdf>

De la Torre AJ, Luat AF, Juhász C, Ho ML, Argersinger DP, Cavuoto KM, Enriquez-Algeciras M, Tikkanen S, North P, Burkhart CN, Chugani HT, Ball KL, Pinto AL, Loeb JA (2018) A Multidisciplinary

Consensus for Clinical Care and Research Needs for Sturge-Weber Syndrome. *Pediatr Neurol.* Jul;84:11-20. doi: 10.1016/j.pediatrneurol.2018.04.005.  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6317878/pdf/nihms961373.pdf>

Ewen JB, Comi AM, Kossoff EH (2007) Myoclonic-astatic epilepsy in a child with Sturge-Weber syndrome. *Pediatr Neurol.* 2007 Feb;36(2):115-7. doi: 10.1016/j.pediatrneurol.2006.08.006.

Gittins S, Steel D, Brunklaus A, Newsom-Davis I, Hawkins C, Aylett SE (2018) Autism spectrum disorder, social communication difficulties, and developmental comorbidities in Sturge-Weber syndrome. *Epilepsy Behav.* 2018 Nov;88:1-4. doi: 10.1016/j.yebeh.2018.08.006.

Hu J, Yu Y, Juhasz C, Kou Z, Xuan Y, Latif Y, Kudo K, Chugani HT, Haacke EM (2008) MR susceptibility weighted imaging (SWI) complements conventional contrast enhanced T1 weighted MRI in characterizing brain abnormalities of Sturge-Weber Syndrome. *J Magn Reson Imaging.* Aug;28(2):300-7. doi: 10.1002/jmri.21435

Jansen FE, Van Der Worp HB, Van Huffelen A, Van Nieuwenhuizen O (2007) Sturge-Weber syndrome and paroxysmal hemiparesis: epilepsy or ischaemia? *Developmental Medicine and Child Neurology.* 46: 783-786 783. doi: 10.1111/j.1469-8749.2004.tb01000.x.  
<https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1469-8749.2004.tb01000.x>

Kossoff EH, Ferenc L, Comi AM (2009) An infantile-onset, severe, yet sporadic seizure pattern is common in Sturge-Weber syndrome. *Epilepsia.* Sep;50(9):2154-7. doi: 10.1111/j.1528-1167.2009.02072.x. Epub 2009 Apr 6. <https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1528-1167.2009.02072.x>

Kossoff EH, Bachur CD, Quain AM, Ewen JB, Comi AM (2014) EEG evolution in Sturge-Weber syndrome. *Epilepsy Res.* May;108(4):816-9. doi: 10.1016/j.eplepsyres.2014.01.023.  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4114141/pdf/nihms603071.pdf>

Kossoff EH, Buck C, Freeman JM (2002) Outcomes of 32 hemispherectomies for Sturge-Weber syndrome worldwide. *Neurology.* Dec 10;59(11):1735-8. doi: 10.1212/01.wnl.0000035639.54567.5c.

Kramer U, Kahana E, Shorer Z, Ben-Zeev B (2000) Outcome of infants with unilateral Sturge-Weber syndrome and early onset seizures. *Dev Med Child Neurol.* Nov;42(11):756-9. doi: 10.1017/s0012162200001407. <https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1469-8749.2000.tb00039.x>

Kwan P, Arzimanoglou A, Berg AT, Brodie MJ, Hauser WA, Mathern G, Moshé SL, Perucca E, Wiebe S, French J (2010) Definition of drug resistant epilepsy: consensus proposal by the ad hoc Task Force of the ILAE Commission on Therapeutic Strategies. *Epilepsia.* Jun;51(6):1069-77. doi: 10.1111/j.1528-1167.2009.02397.x. <https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1528-1167.2009.02397.x>

Lance EI, Lanier KE, Zabel TE, Comi AM (2013) Stimulant use in patients with sturge-weber syndrome: safety and efficacy. *Pediatr Neurol.* Nov;51(5):675-80. doi: 10.1016/j.pediatrneurol.2013.11.009.  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4392725/pdf/nihms-628565.pdf>

Maton B, Krsek P, Jayakar P, Resnick T, Koehn M, Morrison G, Ragheb J, Castellano-Sanchez A, Duchowny M (2010) Medically intractable epilepsy in Sturge-Weber syndrome is associated with cortical malformation: implications for surgical therapy. *Epilepsia.* Feb;51(2):257-67. doi: 10.1111/j.1528-1167.2009.02304.x.

Murakami N, Morioka T, Suzuki SO, Hashiguchi K, Amano T, Sakata A, Iwaki T, Sasaki T (2012) Focal cortical dysplasia type IIa underlying epileptogenesis in patients with epilepsy associated with Sturge-Weber syndrome. *Epilepsia.* Nov;53(11):e184-8. doi: 10.1111/j.1528-1167.2012.03628.x. Epub 2012 Aug 20. <https://onlinelibrary.wiley.com/doi/epdf/10.1111/j.1528-1167.2012.03628.x>

Raches D, Hiscock M, Chapiesski L (2012) Behavioral and academic problems in children with Sturge-Weber syndrome: differences between children with and without seizures. *Epilepsy Behav.* Nov;25(3):457-63. doi: 10.1016/j.yebeh.2012.06.004.

Shirley MD, Tang H, Gallione CJ, Baugher JD, Frelin LP, Cohen B, North PE, Marchuk DA, Comi AM, Pevsner J (2013) Sturge-Weber syndrome and port-wine stains caused by somatic mutation in GNAQ. *N Engl J Med.* May 23;368(21):1971-9. doi: 10.1056/NEJMoa1213507.  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3749068/pdf/nihms491917.pdf>

Sugano H, Nakanishi H, Nakajima M, Higo T, Imura Y, Tanaka Y, Hosozawa M, Niiijima S, Arai H (2014) Posterior quadrant disconnection surgery for Sturge-Weber syndrome. *Epilepsia.* 2014 May;55(5):683-9. doi: 10.1111/epi.12547. <https://onlinelibrary.wiley.com/doi/epdf/10.1111/epi.12547>

Téllez-Zenteno JF, Hernández Ronquillo L, Moien-Afshari F, Wiebe S (2010) Surgical outcomes in lesional and non-lesional epilepsy: a systematic review and meta-analysis. *Epilepsy Res.* May;89(2-3):310-8. doi: 10.1016/j.eplepsyres.2010.02.007.

Tuxhorn IEB, Pannek HW (2002) Epilepsy surgery in bilateral Sturge-Weber syndrome. *Pediatr Neurol.* May;26(5):394-7. doi: 10.1016/s0887-8994(01)00414-3.

Wellmer J, Quesada CM, Rothe L, Elger CE, Bien CG, Urbach H (2013) Proposal for a magnetic resonance imaging protocol for the detection of epileptogenic lesions at early outpatient stages. *Epilepsia.* Nov;54(11):1977-87. doi: 10.1111/epi.12375. Epub 2013 Oct 7.  
<https://onlinelibrary.wiley.com/doi/epdf/10.1111/epi.12375>